



# EFFECT OF THE CLADODES PEELING IN THE FUNCTIONAL, TECHNOLOGICAL, CHEMICAL CHARACTERISTICS AND BIOACTIVE COMPOUNDS IN CLADODES POWDER (*Opuntia ficus-indica*)

SEPULVEDA, E., CATALAN, E., GORENA, T., CHIFFELLE, I. and SAENZ, C.  
 Depto. de Agroindustria y Enología. Facultad de Ciencias Agronómicas, Universidad de Chile. Casilla 1004.  
 Santiago, Chile. e-mail: csaenz@uchile.cl

## INTRODUCTION

Cactus cladodes powder is rich in dietary fiber and could be used as a functional ingredient in foods. The herbaceous aroma and the relation soluble/insoluble fiber could be improved preparing the powder using cladodes with or without epidermis.

## OBJECTIVE

To study the functional, technological and chemical characteristics of 2 types of cactus cladodes powder (with and without epidermis).

## MATERIALS and METHODS

Cladodes (2-3 years) from the University of Chile Garden varieties were used. The cladodes were washed, sliced (0.5 cm) and dried in an air forced tunnel (80°C, 6 hours). Two grounds (blades mills) were realized until a particle size close to 200 $\mu$ . Apparent density (g mL<sup>-1</sup>), moisture (g 100g<sup>-1</sup>), water retention capacity (WRC), swelling (CH) (g mL<sup>-1</sup>), water absorption capacity (WAC), oil absorption index (OAC) (Femenia et al., 1997), and crude and dietary fiber (SDF and IDF) were determined in the powders (Lee et al., 1992). Antioxidant capacity was measured using the DPPH method (TEAC and EC50) (Brand-Williams et al., 1995). Total phenolic content was determined using the Folin-Ciocalteu reagent (mg GAE 100 g<sup>-1</sup>) (Singleton and Rossi, 1965). The data were analyzed by ANOVA using Statgraphics Extra 5.1 program.

## RESULTS and DISCUSSION

The results showed that there was not differences (p<0,05) in the technological characteristics of the powders for moisture, WRC, CH, WAC, OAC (Table 1).



Table 1. Technological characteristics

Parameter	Whole nopal	Peeled nopal
Apparent density (g mL <sup>-1</sup> )	0.50	0.54
Moisture (g 100g <sup>-1</sup> )	6.77 ± 0.19	5.93 ± 0.22
Dry matter (g 100g <sup>-1</sup> )	93.23 ± 0.19	93.89 ± 0.22
Water absorption capacity (g g <sup>-1</sup> )	8.94 ± 0.52	9.51 ± 0.14
Swelling (g mL <sup>-1</sup> )	5.86 ± 0.09	6.49 ± 0.76
Oil absorption index (g g <sup>-1</sup> )	1.48 ± 0.09	1.34 ± 0.08
Water retention capacity (g mL <sup>-1</sup> )	10.22 ± 0.13	10.08 ± 0.15



There was significant differences for the SDF and for the bioactive composition (Table 2); where the powder from cladodes with epidermis have a greater SDF content (14.03±0.38 g 100g<sup>-1</sup>) similar to that reported by Sáenz et al. (2010) and greater phenolic concentration (4.32±0.06 mg GAE g<sup>-1</sup>), capacity antioxidant (1.86±0.02 TEAC) and lipids (2.74±0.69 g 100 g<sup>-1</sup>) compared to powders obtained from cladodes without epidermis (2.96±0.07 mg GAE g<sup>-1</sup>; 1.03±0.05 TEAC; 0.66±0.73 g 100 g, respectively) (Table 3).

Table 2. Dietary fiber

Parameter	Whole nopal	Peeled nopal
Insoluble fiber (IDF) g 100g <sup>-1</sup>	41,9 ± 2,33	41,66 ± 1,08
Soluble fiber (SDF) g 100g <sup>-1</sup>	14,03 ± 0,38	9,15 ± 0,19

Table 3. Bioactive compounds and antioxidant capacity

Parameter	Whole nopal	Peeled nopal
Total phenolics (mg GAE g <sup>-1</sup> )	4.32 ± 0.06	2.96 ± 0.07
Antioxidant activity EC <sub>50</sub>	0.27 ± 0.02	0.47 ± 0.01
Antioxidant activity (TEAC)	1.86 ± 0.02	1.03 ± 0.05

The ash content and crude fiber were greater (p>0,05), in the powders from cladodes without epidermis (10.59±1.06 g 100g<sup>-1</sup> and 7.25±0.29 g 100 g<sup>-1</sup>, respectively) compared with the whole cladodes powders (0.75±0.28 g 100g<sup>-1</sup> and 6.03±0.02 g 100g<sup>-1</sup>, respectively). This unexpected difference can be attributed to the peeling that it could eliminate part of the epidermis layers containing mucilage (soluble fiber). In conclusion the use of whole cladodes, besides avoiding the peeling, provides powders with greater dietary fiber and bioactive compounds content.

## CONCLUSIONS

The cladodes powder including the peel shows greater phenolic content with higher antioxidant activity. Having also a higher content of soluble fiber than the powder of peeled cladodes, this can be due to the epidermis mucilages.

## References

- Brand-Williams, W., Cuvelier, M. E. and Berset, C., 1995. Use of a free radical method to evaluate antioxidant activity. *Lebensm.-Wiss. Technol.* 239, 70-76
- Lee, S.C., Prosky, L. and Devries, J. 1992. Determination of Total, Soluble and Insoluble Fiber Foods. Enzymatic Gravimetric Method. Mes - Tris Buffer: Collaborative Study. *J. Assoc. - Off. Anal. Chem.* 75 (3) 395 - 416.
- Femenia, A., Lefebvre, C., Thebaudin, Y., Robertson, J.A. and Bourgeois, M. 1997. Physical and Sensory Properties of Models Foods Supplemented with Cauliflower Fiber. *Journal of Food Science* 62 (4): 635 - 639.
- Singleton, V.L., and Rossi, J.A. 1965. Colorimetric of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Amer. J. Enol. Vitic.* 16 : 144-158.
- Sáenz, C., Sepúlveda, E., Pak, N. and Lecaros, M. 2010. Chemical and physical characterization of cactus cladodes (*Opuntia ficus-indica*) powder. *Italian Journal of Food Science* 22:4 (In press)

## Acknowledgement

Our research was supported by a Grant from Innova Corfo-Chile N° 07CT9 Z-04, Project "Ingredientes patentables para la industria alimentaria con potencial funcional en la prevención y control de la obesidad y diabetes".

